

Patient Agreement to Systemic Therapy: Consent Form	
BEACOPP-14	
Patient's details / addressograph:	<input type="checkbox"/> Male <span style="margin-left: 200px;"><input type="checkbox"/> Female</span>
	Special requirements (e.g. other language/other communication method)
	<b>Consultant:</b>

**Name of proposed course of treatment: BEACOPP-14**

Bleomycin; IV infusion on day 8

Etoposide; IV infusion on days 1 to 3

Doxorubicin (also known as **Adriamycin**); IV bolus on day 1

Cyclophosphamide; IV bolus on day 1

Vincristine (also known as **Oncovin**); IV bolus on day 8

Procarbazine; oral on days 1 to 7

Prednisolone: oral on days 1 to 7

Repeated every 2 weeks for up to 6 cycles

Macmillan/CRUK/other leaflet(s) given 

**Statement of health professional** (to be filled in by health professional with appropriate knowledge of proposed procedure, as specified in consent policy)

I have discussed what the treatment is likely to involve (including inpatient / outpatient treatment, timing of the treatment, follow-up appointments) and location.

**The intended benefits**

Curative – to give you the best possible chance of being cured

Disease control - the aim is not to cure but to control the disease. The aim is to improve both quality of life and survival

**Significant, unavoidable or frequently occurring risks:** as lymphoma cells die, their breakdown products can damage the kidneys (tumour-lysis syndrome). If you are at risk you will be given medicines to prevent this

**Common side-effects:** low white blood cells (neutropenia) with risk of infection, low red blood cells (anaemia), low platelets with risk of bleeding, feeling sick (nausea) and being sick (vomiting), hair loss, sore mouth and ulcers, numbness or tingling in hands or feet, constipation, chills and fever like symptoms during bleomycin infusion, tiredness and feeling weak

Side effects related to prednisolone: irritation of the stomach lining, increased appetite, changes in blood sugar levels, fluid retention, changes in behaviour (mood swings, difficulty sleeping, anxiety or irritability)

**Less common but potentially life-threatening side-effects:** severe infection causing organ failure

**Other less common side-effects:** irritation of the bladder lining, bleomycin can cause damage to the lungs; if you experience symptoms of shortness of breath or a cough, report this to your doctor immediately. Rarely, doxorubicin may cause long-term damage to the heart. Late effects include a very rare chance of a second cancer.

If doxorubicin or vinblastine accidentally leak outside of the vein during administration, it can cause damage to the surrounding tissues, including potential long-term damage to both the function and appearance of the area. This is not common, but if it happens, nursing staff are trained to minimise the risks of longer term damage.

Cancer can increase your risk of developing a blood clot (thrombosis), and having chemotherapy may increase this risk further. A blood clot may cause pain, redness and swelling in a leg, or breathlessness and chest pain - you must tell your doctor straight away if you have any of these symptoms.

Some chemotherapy drugs can damage women's ovaries and men's sperm, with risk of infertility and early menopause in women. I have warned the patient about the likelihood of:

- early menopause in women       infertility (in men and in women)

Some chemotherapy drugs may damage the development of a baby in the womb (foetus), leading to the many risks associated with an abnormal pregnancy. Therefore, I have discussed the issues of protected sex. This is an issue for both men and women. The patient has been advised not to become pregnant/ not to get a partner pregnant during the period of treatment, and for 12 months (women)/ 6 months (men) after treatment has completed

Any other risks: .....

<b>Clinician Signature</b>	
Signed.....	Date .....
Full Name (print) ..... / .....	Job Title.....
(Forename)	(Surname)

**Statement of patient**

<b>Patient Signature</b>	
Please read this form carefully, which describes the benefits and risks of the proposed treatment. You have the right to change your mind at any time, including after you have signed this form.	
<b>I agree</b> to undergo BEACOPP treatment. I understand the treatment and am aware of the potential side-effects arising from this treatment.	
<b>I understand</b> that you cannot give me a guarantee that a particular person will perform the procedure. The person will, however, have appropriate training and experience.	
Signed.....	Name.....
Date.....	
<b>A witness should sign below if the patient is unable to sign but has indicated his or her consent.</b>	
Signature .....	Date .....
Name (PRINT) .....	

**Statement of interpreter** (where appropriate)

I have interpreted the information above to the patient to the best of my ability and in a way in which I believe s/he can understand.

Signed ..... Name (PRINT) ..... Date.....

**Confirmation of consent** (to be completed by the chemotherapy nurse when the patient attends for the first cycle)

On behalf of the team treating the patient, I have confirmed with the patient that s/he has no further questions and wishes the treatment to go ahead.

Signed ..... Name (PRINT) ..... Date.....

**Copy accepted by patient: yes/no (please ring)**  
**Copy to be retained in patient's notes**

Reason for Update: N/A	Checked and approved by Consultant: Dr E Grey-Davies
Version: 1	Date: 14/12/17